

Resumption of antiplatelet therapy in patients with primary intracranial haemorrhage: Balancing benefits and risks

Khan, Ahsan A.; Lip, Gregory Y.h.

DOI:

[10.1016/j.jns.2017.11.038](https://doi.org/10.1016/j.jns.2017.11.038)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Khan, AA & Lip, GYH 2018, 'Resumption of antiplatelet therapy in patients with primary intracranial haemorrhage: Balancing benefits and risks', *Journal of the Neurological Sciences*, vol. 384, pp. 139-140. <https://doi.org/10.1016/j.jns.2017.11.038>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy


While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Resumption of antiplatelet therapy in patients with primary intracranial haemorrhage: Balancing benefits and risks

Ahsan A. Khan, Gregory Y.H. Lip



PII: S0022-510X(17)34478-7
DOI: doi:[10.1016/j.jns.2017.11.038](https://doi.org/10.1016/j.jns.2017.11.038)
Reference: JNS 15684

To appear in: *Journal of the Neurological Sciences*

Received date: 27 November 2017

Accepted date: 29 November 2017

Please cite this article as: Ahsan A. Khan, Gregory Y.H. Lip , Resumption of antiplatelet therapy in patients with primary intracranial haemorrhage: Balancing benefits and risks. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Jns(2017), doi:[10.1016/j.jns.2017.11.038](https://doi.org/10.1016/j.jns.2017.11.038)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

EDITORIAL

**Resumption of Antiplatelet Therapy in Patients with Primary Intracranial Haemorrhage:
Balancing Benefits and Risks**

Ahsan A Khan, MRCP¹, Gregory Y.H. Lip, MD ^{1,2}

¹. Institute of Cardiovascular Sciences, University of Birmingham, United Kingdom

². Aalborg Thrombosis Research Unit, Department of Clinical Medicine, Faculty of Health, Aalborg University, Aalborg, Denmark.

Correspondence to:

Professor Gregory Y H Lip, MD

Phone +44 121 5075080; g.y.h.lip@bham.ac.uk

Disclosures

Professor Lip has served as a consultant for Bayer/Janssen, BMS/Pfizer, Biotronik, Medtronic, Boehringer Ingelheim, Microlife and Daiichi-Sankyo; and a speaker for Bayer, BMS/Pfizer, Medtronic, Boehringer Ingelheim, Microlife, Roche and Daiichi-Sankyo. No personal fees received.

Keywords: Antiplatelets; intracranial haemorrhage; relative risk; ischaemic stroke; thromboembolism

Intracranial or intracerebral haemorrhage (ICH) can be primary ICH (in the absence of a single clear underlying lesion) and secondary ICH (as a complication of a pre-existing lesion, such as vascular malformation or tumour; or antithrombotic therapy). Primary ICH is the most common type of ICH, and overall, is the second most common subtype of stroke accounting for approximately 10 to 20% of all strokes.

Antiplatelet therapies, particularly aspirin has been the cornerstone for primary and secondary prevention of ischaemic events such as myocardial infarction and ischaemic stroke for over 100 years. As the population ages, the prevalence of cardiovascular and cerebrovascular diseases is rising, leading to an increased use of antiplatelet agents. Conditions such as myocardial infarction and ischaemic stroke have a considerable risk of recurrence, thus necessitating continued use of antiplatelet therapy.

As antiplatelet therapy results in reduction of functioning platelets, theoretically it can predispose to bleeding or delay haemostasis to be achieved after a bleeding event. A systematic review by McQuaid *et al* showed that the use of antiplatelet therapy is associated with an increased risk of haemorrhage, including ICH [1]. People with ICH tend to have significant associated morbidity and mortality and due to the perceived risk of ICH recurrence, primary ICH is usually considered as a relative contraindication for antiplatelet therapy. Thus, the majority of clinicians have adopted a cautious approach in starting or resuming antiplatelet therapy in patients with confirmed diagnosis of ICH. This has serious implications for patients on antiplatelet therapy for a cardiovascular or cerebrovascular indication, who develop primary ICH.

Although few cohort studies have taken place to assess the risk of resuming antiplatelet therapy, these have involved small numbers [2-4]. Therefore, the meta-analysis by Ding and colleagues in this issue of *Journal of the Neurological Sciences* is timely. They performed a methodical and thorough search of major databases and found 6 relevant cohort studies with 1916 patients in total. Their results showed a significantly reduced risk of ischaemic or thromboembolic events following resumption of antiplatelet therapy (relative risk, RR 0.61; 95% confidence interval (CI), 0.48 - 0.79; $P < 0.01$), which perhaps comes as no big surprise. More crucially, they found no significant difference in the risk of ICH recurrence or

haematoma expansion between patients with or without antiplatelet therapy resumption (RR 0.84; 95% CI, 0.47 - 1.51; P=0.56).

These findings are similar to the study by Risselada *et al* which provided evidence that the use of antiplatelets was not associated with an increased risk of ICH [5]. Furthermore, a systematic review performed by Hawryluk *et al* looking at resumption of anticoagulation following central nervous system (CNS) haemorrhage found that early re-initiation of anticoagulants may be prudent [6]. Another meta-analysis of controlled studies by Paciaroni *et al* looking at efficacy and safety of anticoagulants for the prevention of venous thromboembolism in patients with acute haemorrhagic stroke found that early anticoagulation is associated with a significant reduction in PE and a non-significant increase in haematoma enlargement [7].

This meta-analysis by Ding *et al* has several positive aspects addressing an important and clinically relevant issue. Nevertheless, limitations of this meta-analysis should be recognised. The cohort studies did not include patients on dual antiplatelet therapy, which is widely used in the context of primary coronary intervention (PCI) for myocardial infarction and shown to have increased risk of ICH [8, 9]. Only 6 cohort studies were included, and the limited studies and numbers of patients (n=1916) may decrease the reliability of the results. Furthermore, this may underestimate the risk of ICH recurrence or haematoma expansion.

The included studies used different antiplatelet agents, each with a varying bleeding profile. Indeed, 2 of the 6 studies' results showed heterogeneity when assessing the relationship between antiplatelet resumption and risk of ICH recurrence or haematoma expansion [10, 11]. In one of these studies, only aspirin was used which is associated with higher risk of adverse effects including haemorrhagic events when compared with other antiplatelet agents [10]. One other study conducted on patients with atrial fibrillation (AF) found that patients with AF who resumed antiplatelet therapy had decreased risk of ICH recurrence compared to those who did not resume [11]. However, baseline risks of haemorrhagic event between the two groups in the AF study was unknown, and it is possible that the patients with antiplatelet resumption may have had a lower risk of ICH recurrence at baseline due to potential selection bias. Also differences in timing of antiplatelet resumption, location of ICH

and the duration of follow-up and other factors may contribute to the heterogeneity. In addition, the studies included were all observational studies and patients may have difficulty in maintaining adequate adherence to the antiplatelet therapy. Lastly, differences in demographics, ethnicity, lifestyle and comorbidities cannot be avoided in cohort studies leading to potential bias.

What are the future considerations from this work? The meta-analysis by Ding and colleagues has shown that resuming antiplatelet therapy is generally safe in the setting of primary ICH with no significant increase in risk of recurrence or haematoma expansion, whilst providing significant reduction in the risk of ischaemic and thromboembolic events. Clearly, there are several limitations thus warranting a need for well-designed randomised controlled trials to provide better evidence. One such trial, REstart or STop Antithrombotics Randomised Trial (RESTART) is currently underway looking at this particular therapeutic dilemma. The data completion is set to complete by May 2018 and will provide stronger evidence.

For the time being, each individual patient's risks and benefits should be evaluated to enable appropriate management of these patients. Accurate assessment of bleeding risks (and addressing modifiable bleeding risk factors e.g. uncontrolled blood pressure, etc) may help clinicians to identify those patients for whom the benefits of antiplatelet therapy outweigh the risks. A limited number of prediction models are available that predict ICH or major bleeding in patients on antiplatelet therapy [12]. Unfortunately, external validation showed poor performance and thus none can be recommended for use in clinical practice [12]. Biomarkers may offer some insights, but would need to be balanced against the simplicity and practicality of clinical scores [13, 14]. Thus, there is still a great need to develop clinically validated risk scores according to current standards to help guide clinical management, such as those used successfully in the setting of AF and anticoagulation for stroke prophylaxis (e.g. CHA₂DS₂VASc and HAS-BLED risk scores).

References

- [1] K.R. McQuaid, L. Laine, Systematic review and meta-analysis of adverse events of low-dose aspirin and clopidogrel in randomized controlled trials, *Am. J. Med.*, 119 (2006) 624-638.
- [2] R.W.V. Flynn, T.M. MacDonald, G.D. Murray, R.S. MacWalter, A.S.F. Doney, Prescribing Antiplatelet Medicine and Subsequent Events After Intracerebral Hemorrhage, *Stroke*, 41 (2010) 2606-2611.
- [3] A. Biffi, A. Halpin, A. Towfighi, A. Gilson, K. Busl, N. Rost, E.E. Smith, M.S. Greenberg, J. Rosand, A. Viswanathan, Aspirin and recurrent intracerebral hemorrhage in cerebral amyloid angiopathy, *Neurology*, 75 (2010) 693-698.
- [4] B.H. Chong, K.H. Chan, V. Pong, K.K. Lau, Y.H. Chan, M.L. Zuo, W.M. Lui, G.K. Leung, C.P. Lau, H.F. Tse, J.K. Pu, C.W. Siu, Use of aspirin in Chinese after recovery from primary intracranial haemorrhage, *Thromb. Haemost.*, 107 (2012) 241-247.
- [5] R. Risselada, H. Straatman, F. Van Kooten, D.W.J. Dippel, A. Van Der Lugt, W.J. Niessen, A. Firouzan, R.M.C. Herings, M.C.J.M. Sturkenboom, Platelet aggregation inhibitors, vitamin K antagonists and risk of subarachnoid hemorrhage, *J. Thromb. Haemost.*, 9 (2011) 517-523.
- [6] G.W.J. Hawryluk, J.W. Austin, J.C. Furlan, J.B. Lee, C. O'Kelly, M.G. Fehlings, Management of anticoagulation following central nervous system hemorrhage in patients with high thromboembolic risk, *J. Thromb. Haemost.*, 8 (2010) 1500-1508.
- [7] M. Paciaroni, G. Agnelli, M. Venti, A. Alberti, M. Acciarresi, V. Caso, Efficacy and safety of anticoagulants in the prevention of venous thromboembolism in patients with acute cerebral hemorrhage: a meta-analysis of controlled studies, *J. Thromb. Haemost.*, 9 (2011) 893-898.
- [8] S.J. Connolly, J. Pogue, R.G. Hart, S.H. Hohnloser, M. Pfeffer, S. Chrolavicius, S. Yusuf, Effect of clopidogrel added to aspirin in patients with atrial fibrillation, *N. Engl. J. Med.*, 360 (2009) 2066-2078.
- [9] R. Sorensen, M.L. Hansen, S.Z. Abildstrom, A. Hvelplund, C. Andersson, C. Jorgensen, J.K. Madsen, P.R. Hansen, L. Kober, C. Torp-Pedersen, G.H. Gislason, Risk of bleeding in patients with acute myocardial infarction treated with different combinations of aspirin, clopidogrel, and vitamin K antagonists in Denmark: a retrospective analysis of nationwide registry data, *Lancet*, 374 (2009) 1967-1974.
- [10] G.X. T. Chen, D. Tan, C. Wu, Effects of platelet infusion, anticoagulant and other risk factors on the rehaemorrhagia after surgery of hypertensive cerebral hemorrhage, *Eur. Rev. Med. Pharmacol. Sci.*, 19 (2015) 795-799.
- [11] P.B. Nielsen, T.B. Larsen, F. Skjoth, A. Gorst-Rasmussen, L.H. Rasmussen, G.Y. Lip, Restarting Anticoagulant Treatment After Intracranial Hemorrhage in Patients With Atrial Fibrillation and the Impact on Recurrent Stroke, Mortality, and Bleeding: A Nationwide Cohort Study, *Circulation*, 132 (2015) 517-525.
- [12] N.A. Hilken, A. Algra, J.P. Greving, Prediction models for intracranial hemorrhage or major bleeding in patients on antiplatelet therapy: a systematic review and external validation study, *J. Thromb. Haemost.*, 14 (2016) 167-174.
- [13] N.A. Zakai, N.C. Olson, S.E. Judd, D.O. Kleindorfer, B.M. Kissela, G. Howard, M. Cushman, Haemostasis biomarkers and risk of intracerebral haemorrhage in the REasons for Geographic and Racial Differences in Stroke Study, *Thromb. Haemost.*, 117 (2017) 1808-1815.

- [14] M.A. Esteve-Pastor, J.M. Rivera-Caravaca, V. Roldan, V. Vicente, M. Valdes, F. Marin, G.Y.H. Lip, Long-term bleeding risk prediction in 'real world' patients with atrial fibrillation: Comparison of the HAS-BLED and ABC-Bleeding risk scores. The Murcia Atrial Fibrillation Project, *Thromb. Haemost.*, 117 (2017) 1848-1858.